

MAC 1105  
Module 1  
Introduction to Functions and  
Graphs

---

---

---

---

---

---

---

---

**Learning Objectives**

Upon completing this module, you should be able to:

1. Recognize common sets of numbers.
2. Understand scientific notation and use it in applications.
3. Find the domain and range of a relation.
4. Graph a relation in the  $xy$ -plane.
5. Understand function notation.
6. Define a function formally.
7. Identify the domain and range of a function.
8. Identify functions.
9. Identify and use constant and linear functions.
10. Interpret slope as a rate of change.

Rev.S08 <http://faculty.valenciacollege.edu/ashaw/> 2  
Click link to download other modules.

---

---

---

---

---

---

---

---

**Learning Objectives**

11. Write the point-slope and slope-intercept forms for a line.
12. Find the intercepts of a line.
13. Write equations for horizontal, vertical, parallel, and perpendicular lines.
14. Write equations in standard form.
15. Identify and use nonlinear functions.
16. Recognize linear and nonlinear data.
17. Use and interpret average rate of change.
18. Calculate the difference quotient.

Rev.S08 <http://faculty.valenciacollege.edu/ashaw/> 3  
Click link to download other modules.

---

---

---

---

---

---

---

---

## Introduction to Functions and Graphs

There are four major topics in this module:

- Functions and Models
- Graphs of Functions
- Linear Functions
- Equations of Lines

Rev.S08

<http://faculty.valenciacollege.edu/ashaw/>  
Click link to download other modules.

4

---

---

---

---

---

---

---

---

Let's get started by recognizing some common set of numbers.

Rev.S08

<http://faculty.valenciacollege.edu/ashaw/>  
Click link to download other modules.

5

---

---

---

---

---

---

---

---

What is the difference between Natural Numbers and Integers?

• **Natural Numbers** (or counting numbers) are numbers in the set  $N = \{1, 2, 3, \dots\}$ .

• **Integers** are numbers in the set  $I = \{\dots -3, -2, -1, 0, 1, 2, 3, \dots\}$ .

Rev.S08

<http://faculty.valenciacollege.edu/ashaw/>  
Click link to download other modules.

6

---

---

---

---

---

---

---

---

## What are Rational Numbers?

**Rational Numbers** are real numbers which can be expressed as the ratio of two integers  $p/q$  where  $q \neq 0$

**Examples:**  $3 = 3/1$   $-5 = -10/2$   $0 = 0/2$

$0.5 = 1/2$   $0.52 = 52/100$   $0.333... = 1/3$

**Note that:**

- Every integer is a rational number.
- Rational numbers can be expressed as decimals which either terminate (end) or repeat a sequence of digits.

Rev.S08

<http://faculty.valenciacollege.edu/jashaw/>  
Click link to download other modules.

7

---

---

---

---

---

---

---

---

## What are Irrational Numbers?

- **Irrational Numbers** are real numbers which are not rational numbers.
- Irrational numbers Cannot be expressed as the ratio of two integers.
- Have a decimal representation which does not terminate and does not repeat a sequence of digits.

**Examples:**

$\sqrt{2}$ ,  $\sqrt[3]{5}$ ,  $\pi$ ,  $0.01001000100001...$

Rev.S08

<http://faculty.valenciacollege.edu/jashaw/>  
Click link to download other modules.

8

---

---

---

---

---

---

---

---

## Classifying Real Numbers

Classify each number as one or more of the following:  
natural number, integer, rational number, irrational number.

$\sqrt{25}$ ,  $\sqrt[3]{8}$ ,  $3.14$ ,  $.01010101...$ ,  $\frac{22}{7}$ ,  $-\sqrt{11}$

$\sqrt{25} = 5$  so it is a natural number, integer, rational number

$\sqrt[3]{8} = 2$  so it is a natural number, integer, rational number

$3.14$ ,  $.01010101...$ , and  $\frac{22}{7}$  are rational numbers.

$-\sqrt{11}$  is an irrational number.

Rev.S08

<http://faculty.valenciacollege.edu/jashaw/>  
Click link to download other modules.

9

---

---

---

---

---

---

---

---

## Let's Look at Scientific Notation

A real number  $r$  is in **scientific notation** when  $r$  is written as  $c \times 10^n$ , where  $c$  is a real number and  $n$  is an integer.

### Examples:

The distance to the sun is 93,000,000 mi.  
In scientific notation for this is  $9.3 \times 10^7$  mi.

The size of a typical virus is .000005 cm.  
In scientific notation for this is  $5 \times 10^{-6}$  cm.

Rev.S08

<http://faculty.valenciacollege.edu/ashaw/>  
Click link to download other modules.

10

---

---

---

---

---

---

---

---

## What is a Relation? What are Domain and Range?

A **relation** is a set of ordered pairs.

If we denote the ordered pairs by  $(x, y)$

The set of all  $x$  - values is the **DOMAIN**.

The set of all  $y$  - values is the **RANGE**.

### Example

The **relation**  $\{(1, 2), (-2, 3), (-4, -4), (1, -2), (-3, 0), (0, -3)\}$   
has **domain**  $D = \{-4, -3, -2, 0, 1\}$   
and **range**  $R = \{-4, -3, -2, 0, 2, 3\}$

Rev.S08

<http://faculty.valenciacollege.edu/ashaw/>  
Click link to download other modules.

11

---

---

---

---

---


---

---

---

## How to Represent a Relation in a Graph?

The **relation**  $\{(1, 2), (-2, 3), (-4, -4), (1, -2), (-3, 0), (0, -3)\}$   
has the following graph:



Rev.S08

<http://faculty.valenciacollege.edu/ashaw/>  
Click link to download other modules.

12

---

---

---

---

---

---

---

---

## Is Function a Relation?

Recall that a relation is a set of ordered pairs  $(x,y)$  .  
If we think of values of  $x$  as being inputs and values of  $y$  as being outputs, a function is a relation such that for each input there is exactly one output.

This is symbolized by  $output = f(input)$  or  
 $y = f(x)$

Rev.S08

<http://faculty.valenciacollege.edu/ashaw/>  
Click link to download other modules.

13

---

---

---

---

---

---

---

---

## Function Notation

$y = f(x)$

- Is pronounced "y is a function of x."
- Means that given a value of  $x$  (input), there is exactly one corresponding value of  $y$  (output).
- $x$  is called the independent variable as it represents inputs, and  $y$  is called the dependent variable as it represents outputs.
- Note that:  $f(x)$  is NOT  $f$  multiplied by  $x$ .  $f$  is NOT a variable, but the name of a function (the name of a relationship between variables).

Rev.S08

<http://faculty.valenciacollege.edu/ashaw/>  
Click link to download other modules.

14

---

---

---

---

---

---

---

---

## What are Domain and Range?

The set of all meaningful inputs is called the **DOMAIN** of the function.

The set of corresponding outputs is called the **RANGE** of the function.

Rev.S08

<http://faculty.valenciacollege.edu/ashaw/>  
Click link to download other modules.

15

---

---

---

---

---

---

---

---

## What is a Function?

A **function** is a relation in which each element of the domain corresponds to exactly one element in the range.

The function may be defined by a set of ordered pairs, a table, a graph, or an equation.

Rev.S08

<http://faculty.valenciacollege.edu/ashaw/>  
Click link to download other modules.

16

---

---

---

---

---

---

---

---

## Here is an Example

- Suppose a car travels at 70 miles per hour. Let  $y$  be the distance the car travels in  $x$  hours. Then  $y = 70x$ .
- Since for each value of  $x$  (that is the time in hours the car travels) there is just one corresponding value of  $y$  (that is the distance traveled),  $y$  is a function of  $x$  and we write
$$y = f(x) = 70x$$
- Evaluate  $f(3)$  and interpret.
  - $f(3) = 70(3) = 210$ . This means that the car travels 210 miles in 3 hours.

Rev.S08

<http://faculty.valenciacollege.edu/ashaw/>  
Click link to download other modules.

17

---

---

---

---

---

---

---

---

## Here is Another Example

- Given the following data, is  $y$  a function of  $x$ ?

– Input	$x$	3	4	8
– Output	$y$	6	6	-5
- Note: The data in the table can be written as the set of ordered pairs  $\{(3,6), (4,6), (8,-5)\}$ .
- Yes,  $y$  is a function of  $x$ , because for each value of  $x$ , there is just one corresponding value of  $y$ . Using function notation we write  $f(3) = 6$ ;  $f(4) = 6$ ;  $f(8) = -5$ .

Rev.S08

<http://faculty.valenciacollege.edu/ashaw/>  
Click link to download other modules.

18

---

---

---

---

---

---

---

---

## One More Example

- Undergraduate Classification at Study-Hard University (SHU) is a function of Hours Earned. We can write this in function notation as  $C = f(H)$ .
- Why is  $C$  a function of  $H$ ?
  - For each value of  $H$  there is exactly one corresponding value of  $C$ .
  - In other words, for each input there is exactly one corresponding output.

Rev.S08

<http://faculty.valenciacollege.edu/ashaw/>  
Click link to download other modules.

19

---

---

---

---

---

---

---

---

## One More Example (Cont.)

- Here is the classification of students at SHU (from catalogue):
- No student may be classified as a sophomore until after earning at least 30 semester hours.
- No student may be classified as a junior until after earning at least 60 hours.
- No student may be classified as a senior until after earning at least 90 hours.

Rev.S08

<http://faculty.valenciacollege.edu/ashaw/>  
Click link to download other modules.

20

---

---

---

---

---

---

---

---

## One More Example (Cont.)

- Remember  $C = f(H)$
- Evaluate  $f(20)$ ,  $f(30)$ ,  $f(0)$ ,  $f(20)$  and  $f(61)$ :
  - $f(20) = \text{Freshman}$
  - $f(30) = \text{Sophomore}$
  - $f(0) = \text{Freshman}$
  - $f(61) = \text{Junior}$
- What is the domain of  $f$ ?
- What is the range of  $f$ ?

Rev.S08

<http://faculty.valenciacollege.edu/ashaw/>  
Click link to download other modules.

21

---

---

---

---

---

---

---

---

## One More Example (Cont.)

**Domain** of  $f$  is the set of **non-negative integers**  $\{0, 1, 2, 3, 4, \dots\}$

Alternatively, some individuals say the domain is the set of **positive rational numbers**, since technically one could earn a fractional number of hours if they transferred in some quarter hours. For example, 4 quarter hours =  $2 \frac{2}{3}$  semester hours.

Some might say the domain is the set of **non-negative real numbers**  $[0, \infty)$ , but this set includes irrational numbers. It is impossible to earn an irrational number of credit hours. For example, one could not earn  $\sqrt{2}$  hours.

**Range** of  $f$  is  $\{\text{Fr, Soph, Jr, Sr}\}$

Rev.S08

<http://faculty.valenciacollege.edu/ashaw/>  
Click link to download other modules.

22

---

---

---

---

---

---

---

---

## Identifying Functions

Referring to the previous example concerning SHU, is hours earned a **function of classification**? That is, is  $H = f(C)$ ? Explain why or why not.

Is classification a **function** of years spent at SHU? Why or why not?

- Given  $x = y^2$ , is  $y$  a **function of  $x$** ? Why or why not?
- Given  $x = y^2$ , is  $x$  a **function of  $y$** ? Why or why not?
- Given  $y = x^2 + 7$ , is  $y$  a **function of  $x$** ? Why, why not?

Rev.S08

<http://faculty.valenciacollege.edu/ashaw/>  
Click link to download other modules.

23

---

---

---

---

---

---

---

---

## Identifying Functions (Cont.)

- Is hours earned a function of classification? That is, is  $H = f(C)$ ?
- That is, for each value of  $C$  is there just one corresponding value of  $H$ ?
  - No. One example is
    - if  $C = \text{Freshman}$ , then  $H$  could be 3 or 10 (or lots of other values for that matter)

Rev.S08

<http://faculty.valenciacollege.edu/ashaw/>  
Click link to download other modules.

24

---

---

---

---

---

---

---

---



### Identifying Functions (Cont.)

- Is classification a function of years spent at SHU? That is, is  $C = f(Y)$ ?
- That is, for each value of  $Y$  is there just one corresponding value of  $C$ ?
  - No. One example is
    - if  $Y = 4$ , then  $C$  could be Sr. or Jr. It could be Jr if a student was a part time student and full loads were not taken.

Rev.S08 <http://faculty.valenciacollege.edu/ashaw/> Click link to download other modules. 25

---

---

---

---

---

---

---

---

### Identifying Functions (Cont.)

- Given  $x = y^2$ , is  $y$  a function of  $x$ ?
- That is, given a value of  $x$ , is there just one corresponding value of  $y$ ?
  - No, if  $x = 4$ , then  $y = 2$  or  $y = -2$ .

Rev.S08 <http://faculty.valenciacollege.edu/ashaw/> Click link to download other modules. 26

---

---

---

---

---

---

---

---

### Identifying Functions (Cont.)

- Given  $x = y^2$ , is  $x$  a function of  $y$ ?
- That is, given a value of  $y$ , is there just one corresponding value of  $x$ ?
  - Yes, given a value of  $y$ , there is just one corresponding value of  $x$ , namely  $y^2$ .

Rev.S08 <http://faculty.valenciacollege.edu/ashaw/> Click link to download other modules. 27

---

---

---

---

---

---

---

---

### Identifying Functions (Cont.)

- Given  $y = x^2 + 7$ , is  $y$  a function of  $x$ ?
- That is, given a value of  $x$ , is there just one corresponding value of  $y$ ?
  - Yes, given a value of  $x$ , there is just one corresponding value of  $y$ , namely  $x^2 + 7$ .

Rev.S08 <http://faculty.valenciacollege.edu/ashaw/> 28  
Click link to download other modules.

---

---

---

---

---

---

---

---

### Five Ways to Represent a Function

- Verbally
- Numerically
- Diagrammatically
- Symbolically
- Graphically

Rev.S08 <http://faculty.valenciacollege.edu/ashaw/> 29  
Click link to download other modules.

---

---

---

---

---

---

---

---

### Verbal Representation

- Referring to the previous example:
  - If you have less than 30 hours, you are a freshman.
  - If you have 30 or more hours, but less than 60 hours, you are a sophomore.
  - If you have 60 or more hours, but less than 90 hours, you are a junior.
  - If you have 90 or more hours, you are a senior.

Rev.S08 <http://faculty.valenciacollege.edu/ashaw/> 30  
Click link to download other modules.

---

---

---

---

---

---

---

---

### Numeric Representation

<i>H</i>	<i>C</i>
0	Freshman
1	Freshman
?	
?	
?	
?	
29	Freshman
30	Sophomore
31	Sophomore
?	
?	
?	
59	Sophomore
60	Junior
61	Junior
?	
?	
?	
89	Junior
90	Senior
91	Senior
?	
?	
?	

Rev.S08

<http://faculty.valenciaccc.edu/jshaw/>  
Click link to download other modules.

31

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

### Symbolic Representation

$$C = f(H) = \begin{cases} \text{Freshman} & \text{if } 0 \leq H < 30 \\ \text{Sopho} & \text{if } 30 \leq H < 60 \\ \text{Junior} & \text{if } 60 \leq H < 90 \\ \text{Senior} & \text{if } H \geq 90 \end{cases}$$

Rev.S08

<http://faculty.valenciaccc.edu/jshaw/>  
Click link to download other modules.

32

---

---

---

---

---

---

---

---

---

---

---


---

---

---

---

---



Rev.S08

<http://faculty.valenciaccc.edu/jshaw/>  
Click link to download other modules.

33

---

---

---

---

---

---

---

---

---

---

---

---

---


---

---

---

## Graphical Representation

- In this graph the domain is considered to be  $[0, \infty)$
- instead of  $\{0, 1, 2, 3, \dots\}$ , and note that inputs are typically graphed on the horizontal axis and outputs are typically graphed on the vertical axis.



Rev.S08

<http://faculty.valenciacollege.edu/jashaw/>  
Click link to download other modules.

34

---

---

---

---

---

---

---

---

## Vertical Line Test

- Another way to determine if a graph represents a function, simply visualize vertical lines in the  $xy$ -plane. If each vertical line intersects a graph at no more than one point, then it is the graph of a function.

Rev.S08

<http://faculty.valenciacollege.edu/jashaw/>  
Click link to download other modules.

35

---

---

---

---

---

---

---

---

## What is a Constant Function?


A function  $f$  represented by  $f(x) = b$ , where  $b$  is a constant (fixed number), is a constant function.

Examples:

$$f(x) = 2$$

$$f(x) = \frac{-1}{2}$$

$$f(x) = \sqrt{2}$$



Note: Graph of a constant function is a horizontal line.

Rev.S08

<http://faculty.valenciacollege.edu/jashaw/>  
Click link to download other modules.

36

---

---

---

---

---

---

---

---

## What is a Linear Function?

A function  $f$  represented by  $f(x) = ax + b$ ,

where  $a$  and  $b$  are constants, is a linear function.


(It will be an identity function, if constant  $a = 1$  and constant  $b = 0$ .)

**Examples:**

$f(x) = 2x + 3$  (Note:  $a = 2$  and  $b = 3$ )

$f(x) = -5x - \frac{1}{2}$  (Note:  $a = -5$  and  $b = -\frac{1}{2}$ )

$f(x) = 2$  (Note:  $a = 0$  and  $b = 2$ )



Note that a  $f(x) = 2$  is both a linear function and a constant function. A constant function is a special case of a linear function.

Rev.S08

<http://faculty.valenciacollege.edu/ashaw/>  
Click link to download other modules.

37

---

---

---

---

---

---

---

---

## Rate of Change of a Linear Function

x	y
-2	-1
-1	1
0	3
1	5
2	7
3	9

Table of values for  $f(x) = 2x + 3$ .

- Note throughout the table, as  $x$  increases by 1 unit,  $y$  increases by 2 units. In other words, the **RATE OF CHANGE** of  $y$  with respect to  $x$  is constantly 2 throughout the table. Since the rate of change of  $y$  with respect to  $x$  is constant, the function is **LINEAR**. Another name for rate of change of a linear function is **SLOPE**.

Rev.S08

<http://faculty.valenciacollege.edu/ashaw/>  
Click link to download other modules.

38

---

---

---

---

---


---

---

---

## The Slope of a Line

- The slope  $m$  of the line passing through the points  $(x_1, y_1)$  and  $(x_2, y_2)$  is



Rev.S08

<http://faculty.valenciacollege.edu/ashaw/>  
Click link to download other modules.

39

---

---

---

---

---

---


---

---

## Example of Calculation of Slope

- Find the **slope** of the line passing through the points  $(-2, -1)$  and  $(3, 9)$ .

$$m = \frac{\Delta y}{\Delta x} = \frac{9 - (-1)}{3 - (-2)} = \frac{10}{5} = 2$$



- The **slope** being 2 means that for each unit  $x$  increases, the corresponding increase in  $y$  is 2. The **rate of change** of  $y$  with respect to  $x$  is 2/1 or 2.

Rev.S08

<http://faculty.valenciacollege.edu/ashaw/>  
Click link to download other modules.

40

---

---

---

---

---

---

---

---

## How to Write the Point-Slope Form of the Equation of a Line?

The line with **slope**  $m$  passing through the **point**  $(x_1, y_1)$  has equation

$$y = m(x - x_1) + y_1$$

or

$$y - y_1 = m(x - x_1)$$

Rev.S08

<http://faculty.valenciacollege.edu/ashaw/>  
Click link to download other modules.

41

---

---

---

---

---

---

---

---

## How to Write the Equation of the Line Passing Through the Points $(-4, 2)$ and $(3, -5)$ ?

To write the equation of the line using **point-slope form**

$$y = m(x - x_1) + y_1$$

the **slope**  $m$  and a **point**  $(x_1, y_1)$  are needed.

Let  $(x_1, y_1) = (3, -5)$ .

Calculate  $m$  using the two given points.

$$m = \frac{\Delta y}{\Delta x} = \frac{-5 - 2}{3 - (-4)} = \frac{-7}{7} = -1$$

Equation is  $y = -1(x - 3) + (-5)$

This simplifies to  $y = -x + 3 + (-5)$

$$y = -x - 2$$

Rev.S08

<http://faculty.valenciacollege.edu/ashaw/>  
Click link to download other modules.

42

---

---

---

---

---

---

---

---

## Slope-Intercept Form

The line with slope  $m$  and  $y$ -intercept  $b$  is given by

$$- y = m x + b$$

Rev.S08

<http://faculty.valenciacollege.edu/ashaw/>  
Click link to download other modules.

43

---

---

---

---

---

---

---

---

## How to Write the Equation of a line passing through the point (0,-2) with slope $\frac{1}{2}$ ?


Since the point (0, -2) has an  $x$ -coordinate of 0, the point is a  $y$ -intercept. Thus  $b = -2$

Using slope-intercept form

$$y = m x + b$$

the equation is

$$y = (\frac{1}{2}) x - 2$$



Rev.S08

<http://faculty.valenciacollege.edu/ashaw/>  
Click link to download other modules.

44

---

---

---

---

---


---

---

---

## How to Write an Equation of a Linear Function in Slope-Intercept Form?

- What is the slope?
  - As  $x$  increases by 4 units,  $y$  decreases by 3 units so the slope is  $-\frac{3}{4}$
- What is the  $y$ -intercept?
  - The graph crosses the  $y$ -axis at (0,3) so the  $y$ -intercept is 3.
- What is the equation?
  - Equation is
  - $f(x) = (-\frac{3}{4})x + 3$



Rev.S08

<http://faculty.valenciacollege.edu/ashaw/>  
Click link to download other modules.

45

---

---

---

---

---

---

---

---

## What is the Standard Form for the Equation of a Line?

$$ax + by = c$$

is **standard form** (or **general form**) for the equation of a line.

Rev.S08

<http://faculty.valenciacollege.edu/ashaw/>  
Click link to download other modules.

46

---

---

---

---

---


---

---

---

## How to Find x-Intercept and y-intercept?

- To find the x-intercept, let  $y = 0$  and solve for  $x$ .
  - $2x - 3(0) = 6$
  - $2x = 6$
  - $x = 3$
- To find the y-intercept, let  $x = 0$  and solve for  $y$ .
  - $2(0) - 3y = 6$
  - $-3y = 6$
  - $y = -2$



Rev.S08

<http://faculty.valenciacollege.edu/ashaw/>  
Click link to download other modules.

47

---

---

---

---

---

---

---

---

## What are the Characteristics of Horizontal Lines?


Slope is 0, since  $\Delta y = 0$  and  $m = \Delta y / \Delta x$

Equation is:  $y = mx + b$

$$y = (0)x + b$$

$y = b$  where  $b$  is the y-intercept

Example:  $y = 3$  (or  $0x + y = 3$ )



Note that regardless of the value of  $x$ , the value of  $y$  is always 3.

Rev.S08

<http://faculty.valenciacollege.edu/ashaw/>  
Click link to download other modules.

48

---

---

---

---

---

---


---

---



## What are the Characteristics of Vertical Lines?

- Slope is undefined, since  $\Delta x = 0$  and  $m = \Delta y / \Delta x$
- Example:



- Note that regardless of the value of  $y$ , the value of  $x$  is always 3.
- Equation is  $x = 3$  (or  $x + 0y = 3$ )
- Equation of a vertical line is  $x = k$  where  $k$  is the  $x$ -intercept.

Rev.S08

<http://faculty.valenciacollege.edu/ashaw/>  
Click link to download other modules.

49

---

---

---

---

---

---

---



---

---

---

## What Are the Differences Between Parallel and Perpendicular Lines?

- Parallel lines have the same slant, thus they have the same slopes.
- Perpendicular lines have slopes which are negative reciprocals (unless one line is vertical!)



Rev.S08

<http://faculty.valenciacollege.edu/ashaw/>  
Click link to download other modules.

50

---

---

---

---

---

---

---

---

---

---


## How to Find the Equation of the Line Perpendicular to $y = -4x - 2$ Through the Point $(3, -1)$ ?

The slope of any line perpendicular to  $y = -4x - 2$  is  $\frac{1}{4}$  ( $-4$  and  $\frac{1}{4}$  are negative reciprocals)

Since we know the slope of the line and a point on the line we can use point-slope form of the equation of a line:

$$y = m(x - x_1) + y_1$$

$$y = (1/4)(x - 3) + (-1)$$



In slope-intercept form:

$$y = (1/4)x - (3/4) + (-1)$$

$$y = (1/4)x - 7/4$$

Rev.S08

<http://faculty.valenciacollege.edu/ashaw/>  
Click link to download other modules.

51

---

---

---

---

---

---

---

---


---

---

### Example of a Linear Function

The table and corresponding graph show the price  $y$  of  $x$  tons of landscape rock.

$x$ (tons)	$y$ (price in dollars)
25	5
75	100



$y$  is a linear function of  $x$  and the slope is  $\frac{\Delta y}{\Delta x} = \frac{100 - 5}{75 - 25} = 25$

The **rate of change** of price  $y$  with respect to tonnage  $x$  is 25 to 1. This means that for an increase of 1 ton of rock the price increases by \$25.

Rev.S08

<http://faculty.valenciacollege.edu/ashaw/>  
Click link to download other modules.

52

---

---

---

---

---

---

---

---

---

---

---

---

### Example of a Nonlinear Function

$x$	$y$
0	0
1	1
2	4




Table of values for  $f(x) = x^2$

Note that as  $x$  increases from 0 to 1,  $y$  increases by 1 unit; while as  $x$  increases from 1 to 2,  $y$  increases by 3 units. 1 does not equal 3. This function **does NOT** have a **CONSTANT RATE OF CHANGE** of  $y$  with respect to  $x$ , so the function is **NOT LINEAR**.

Note that the graph is **not a line**.

Rev.S08

<http://faculty.valenciacollege.edu/ashaw/>  
Click link to download other modules.

53

---

---

---

---

---

---

---

---

---

---

---

---

### Average Rate of Change

Let  $(x_1, y_1)$  and  $(x_2, y_2)$  be distinct points on the graph of a function  $f$ . The **average rate of change** of  $f$  from  $x_1$  to  $x_2$  is

$$\frac{y_2 - y_1}{x_2 - x_1}$$

Note that the **average rate of change** of  $f$  from  $x_1$  to  $x_2$  is the **slope** of the line passing through  $(x_1, y_1)$  and  $(x_2, y_2)$

Rev.S08

<http://faculty.valenciacollege.edu/ashaw/>  
Click link to download other modules.

54

---

---

---

---

---

---

---

---

---


---

---

---

## What is the Difference Quotient?

The **difference quotient** of a function  $f$  is an expression of the form  $\frac{f(x+h) - f(x)}{h}$  where  $h$  is not 0.



Note that a **difference quotient** is actually an **average rate of change**.

Rev.S08

<http://faculty.valenciacollege.edu/ashaw/>  
Click link to download other modules.

55

---

---

---

---

---

---

---

---

## What have we learned?

We have learned to:

1. Recognize common sets of numbers.
2. Understand scientific notation and use it in applications.
3. Find the domain and range of a relation.
4. Graph a relation in the  $xy$ -plane.
5. Understand function notation.
6. Define a function formally.
7. Identify the domain and range of a function.
8. Identify functions.
9. Identify and use constant and linear functions.
10. Interpret slope as a rate of change.

Rev.S08

<http://faculty.valenciacollege.edu/ashaw/>  
Click link to download other modules.

56

---

---

---

---

---

---

---

---

## What have we learned? (Cont.)

11. Write the point-slope and slope-intercept forms for a line.
12. Find the intercepts of a line.
13. Write equations for horizontal, vertical, parallel, and perpendicular lines.
14. Write equations in standard form.
15. Identify and use nonlinear functions.
16. Recognize linear and nonlinear data.
17. Use and interpret average rate of change.
18. Calculate the difference quotient.

Rev.S08

<http://faculty.valenciacollege.edu/ashaw/>  
Click link to download other modules.

57

---

---

---

---

---

---

---

---

## Credit

Some of these slides have been adapted/modified in part/whole from the slides of the following textbook:

- Rockswold, Gary, Precalculus with Modeling and Visualization, 3th Edition

Rev.S08

<http://faculty.valenciacollege.edu/jackhaw/>  
Click link to download other modules.

58

---

---

---

---

---

---

---

---